

<h2>Dapagliflozin Tablets</h2> <h3>DAPACOOOL-10 MG</h3> <p>डापाग्लिफ्लो-10 एमजी Tablets</p>				
10 x 10 Tablets				
<p>Composition : Each film coated tablet contains : Dapagliflozin Propanediol Monohydrate 10 mg. equivalent to Dapagliflozin q.s. Excipients Colours : Titanium Dioxide IP & Iron oxide yellow</p>		<p>Storage : Store protected from moisture at a temperature not exceeding 30°C.</p> <p>CAUTION : Not to be sold by retail without the prescription of a Registered Medical Practitioner.</p> <p>Keep the medicine out of reach of children.</p> <p>Manufactured by: Mascot Health Series Pvt. Ltd. At-Plot No. 52, Sec-6A, IIE, Sidcul, Haridwar-249403, Uttarakhand.</p>		
<p>Dosage : As directed by the Physician.</p>				



DAPACOOOL-10 (Dapagliflozin 10 mg Tablet)

Product Overview: DAPACOOOL-10 contains Dapagliflozin, an oral medication used in the management of type 2 diabetes mellitus. It helps lower blood sugar levels by preventing glucose reabsorption in the kidneys and promoting its excretion through urine.

Drug Class:

- **Dapagliflozin:** Sodium-Glucose Co-Transporter 2 (SGLT2) Inhibitor

Indications and Uses: DAPACOOOL-10 is indicated for:

- Management of type 2 diabetes mellitus in adults, as an adjunct to diet and exercise.
- It may be used alone or in combination with other antidiabetic medications.

Dosage and Administration:

- **Adults:** The usual recommended dose is one tablet (10 mg) once daily, taken in the morning, with or without food.

- **Administration:** Swallow the tablet whole with water. Do not crush or chew the tablet.
- **Important:** Follow your healthcare provider's instructions regarding the duration of treatment and dosage adjustments.

Common Side Effects:

- Urinary tract infections
- Increased urination
- Thirst
- Nausea

Less Common Side Effects:

- Yeast infections
- Low blood pressure (hypotension)
- Dehydration
- Ketoacidosis (a serious condition characterized by high levels of ketones in the blood)

Warnings and Precautions:

- Do not exceed the recommended dosage.
- Monitor blood glucose levels regularly.
- Inform your doctor of any other medications or health conditions.
- Use with caution in patients with a history of urinary tract infections or kidney problems.
- Consult your healthcare provider if you experience symptoms of ketoacidosis (such as nausea, vomiting, abdominal pain, or confusion).

Drug Interactions:

- DAPACOOOL-10 may interact with other medications that affect blood glucose levels, diuretics, or medications that impact kidney function. Inform your healthcare provider about all other medications and supplements you are taking.

Contraindications:

- Hypersensitivity to Dapagliflozin or any components of the product.
- Severe renal impairment or end-stage renal disease.
- Type 1 diabetes mellitus.

Dos and Don'ts: Dos:

- Take the medication as prescribed by your healthcare provider.
- Maintain regular follow-ups and blood glucose monitoring.
- Stay hydrated and report any unusual symptoms to your healthcare provider.

Don'ts:

- Do not take more than the prescribed dose.
- Avoid abrupt discontinuation of the medication without consulting your healthcare provider.
- Do not use if you have severe renal impairment without medical advice.

Before Taking DAPACOOOL-10:

- Discuss your complete medical history with your doctor, including any allergies, existing health conditions, and current medications.
- Inform your doctor if you are pregnant, planning to become pregnant, or are breastfeeding.

Disclaimer: This information is provided for educational purposes only and should not be used for self-medication. Always consult your doctor for appropriate medical advice and treatment. Do not alter the dosage prescribed by your healthcare provider.

Quality Product from Econ Healthcare**Contact Information:**

- **Address:** 11, First Floor, Science City Road, Sola, Ahmedabad

- **Phone:** +91 635-6666-700
- **Email:** info@econhealthcare.net
- **Website:** www.econhealthcare.net

All rights reserved to Econ Healthcare