



<p>Rx Cefixime & Ofloxacin Tablets</p> <p>Cefi cool - O</p> <p>10 x 10 Tablets</p>	
--	--

Each film coated tablet contains :
Cefixime I.P.
Eq. to Cefixime Anhydrous 200 mg.
Ofloxacin I.P. 200 mg.
Excipients Q.S.
Colour : Yellow oxide of Iron I.P.
Dose: As directed by the Physician
Storage: Store in a cool dry place,
Protect from Light.
Keep medicine out of reach of children

SCHEDULE 'H1' - PRESCRIPTION DRUG-CAUTION
It is dangerous to take this preparation except in accordance with the medical advice.
Not to be sold by retail without the prescription of a Registered Medical Practitioner.

Precaution : Antacid containing Calcium, Magnesium or Aluminium, Sucralfate & Iron or Multivitamin Containing Zinc should not be taken within two hours period before or after ofloxacin administration.

CEFICOOL-O TAB

Composition: Cefixime 200mg + Ofloxacin 200mg

General Information

CEFICOOL-O TAB is a combination of Cefixime and Ofloxacin, two potent antibiotics that belong to the cephalosporin and fluoroquinolone classes, respectively. This combination is effective in treating a wide range of bacterial infections by inhibiting bacterial cell wall synthesis and DNA replication, leading to the elimination of harmful bacteria.

Class

- **Cefixime:** Third-generation cephalosporin antibiotic.
- **Ofloxacin:** Fluoroquinolone antibiotic.

Indications and Uses

CEFICOOL-O TAB is used to treat various bacterial infections, including:

- Respiratory tract infections.

- Urinary tract infections (UTIs).
 - Gastrointestinal infections.
 - Typhoid fever.
 - Skin and soft tissue infections.
 - Gynecological infections.
-

Common and Less Common Side Effects

- **Common:** Nausea, diarrhea, headache, dizziness, abdominal pain, and loss of appetite.
 - **Less Common:** Allergic reactions (rash, itching), blurred vision, and joint or muscle pain. Contact your doctor if any side effects persist or worsen.
-

Doses and Administration

- Take CEFICOOOL-O TAB as prescribed by your doctor, typically once or twice daily.
 - Can be taken with or without food, though it is better to take it with food to reduce stomach upset.
 - Complete the full course of treatment even if you feel better to prevent bacterial resistance.
-

Warnings and Precautions

- Avoid use in patients with a history of allergic reactions to cephalosporins or fluoroquinolones.
 - Use cautiously in individuals with kidney or liver impairment.
 - May cause dizziness; avoid driving or operating heavy machinery if affected.
 - Avoid prolonged exposure to sunlight as Ofloxacin may increase photosensitivity.
-

Drug Interactions

- Antacids and iron supplements may reduce the absorption of Ofloxacin; avoid taking them together.
 - Can interact with blood thinners, anti-inflammatory drugs, and other antibiotics.
 - Inform your doctor about all the medications you are taking.
-

Contraindications

- Hypersensitivity to Cefixime, Ofloxacin, or any other components of the medication.

- Not recommended during pregnancy or breastfeeding without a doctor's consultation.
-

Dos and Don'ts

- **Do:** Follow your doctor's prescription carefully and maintain hydration during treatment.
 - **Don't:** Stop taking the medication prematurely or take more than the prescribed dose.
-

Before Taking

Inform your doctor if you:

- Have a history of kidney or liver issues.
 - Are pregnant, planning to conceive, or breastfeeding.
 - Are taking any other medications, including over-the-counter drugs or supplements.
-

Disclaimer

This information is intended for educational purposes only and does not replace a doctor's prescription. Always consult your doctor before starting or stopping any medication. Do not change the prescribed dose without medical advice.

Econ Healthcare Pvt Ltd

A quality product from **Econ Healthcare Pvt Ltd**.

Address: 11, First Floor, Science City Road, Sola, Ahmedabad.

Phone: +91 635-6666-700

Email: info@econhealthcare.net

Website: www.econhealthcare.net

All rights reserved to Econ Healthcare.