



Rx
Dapagliflozin, Glimepiride & Metformin Hydrochloride (Extended Release) Tablets

DAPACOOl-GM 1 FORTE
डापाकूल-जीएम 1 फोर्टे Tablets

10 x 10 Tablets

Composition:
Each film coated bilayered tablet contains:
Dapagliflozin Propanediol USP
Eq. to Dapagliflozin 10 mg
Glimepiride IP 1 mg
Metformin Hydrochloride IP 1000 mg
(As Extended release)
Excipients q.s
Colour: Sunset Yellow Lake
Dosage : As directed by the Physician.
Store protected from light & moisture at a temperature not exceeding 30°C.
Keep the medicine out of reach of children.
Tablet to be taken whole and not chewed.

SCHEDULE G PRESCRIPTION DRUG-CAUTION
It is dangerous to use this preparation except under medical supervision.

SCHEDULE H PRESCRIPTION DRUG-CAUTION
Not to be sold to retail without the prescription of a Registered Medical Practitioner.

Manufactured by: Mascot Health Series Pvt. Ltd.
At-Plot No. 52, Sec-6A, IIE, Sidcul,
Haridwar-249403, Uttarakhand.

Brand Name: DAPACOOl-GM 1 FORTE TAB

Composition: Dapagliflozin 10 mg, Glimepiride 1 mg, Metformin 500 mg

General Information

DAPACOOl-GM 1 FORTE TAB is a combination medication designed to manage **Type 2 Diabetes Mellitus** effectively. This triple-drug formula combines:

- **Dapagliflozin**, an SGLT2 inhibitor that helps eliminate excess glucose through urine.
- **Glimepiride**, a sulfonylurea that boosts insulin secretion from pancreatic beta cells.
- **Metformin**, a biguanide that lowers liver glucose production and enhances insulin sensitivity.

Indications and Uses

- Provides glycemic control in adults with Type 2 Diabetes Mellitus.
- Suitable for patients requiring triple therapy to achieve optimal blood sugar levels.

Common and Less Common Side Effects

Common Side Effects:

- Low blood sugar (hypoglycemia)
- Gastrointestinal issues like nausea, diarrhea, or bloating
- Increased frequency of urination

Less Common Side Effects:

- Dizziness, fatigue, or dehydration
 - Rare cases of urinary tract infections
 - Lactic acidosis (a rare but serious side effect of Metformin)
-

Doses and Administration

- The dosage must be prescribed by a doctor based on individual requirements.
 - Administer orally with meals to minimize gastrointestinal side effects.
 - Do not skip meals while taking this medication.
-

Warnings and Precautions

- Regularly monitor blood glucose levels.
 - Avoid alcohol consumption to prevent hypoglycemia or lactic acidosis.
 - Use with caution if you have kidney, liver, or heart conditions.
-

Drug Interactions

- May interact with other anti-diabetic drugs, diuretics, or corticosteroids.
 - Avoid using with medications that may increase the risk of lactic acidosis.
 - Share details of all current medications with your doctor before starting.
-

Contraindications

- Not suitable for patients with severe kidney or liver impairment.
 - Should not be used in Type 1 Diabetes or diabetic ketoacidosis.
 - Avoid if allergic to any of the ingredients in this medication.
-

Dos and Don'ts

Dos:

- Follow dietary and exercise recommendations as advised by your doctor.
- Stay hydrated and maintain consistent monitoring of glucose levels.

Don'ts:

- Do not alter the dosage without consulting a healthcare professional.
 - Avoid skipping doses or stopping the medication abruptly.
-

Before Taking DAPACOOOL-GM 1 FORTE TAB

- Inform your doctor if you are pregnant, planning to conceive, or breastfeeding.
 - Discuss any history of chronic conditions like heart, kidney, or liver issues.
 - Report any unusual symptoms like persistent nausea, rapid breathing, or dizziness immediately.
-

Econ Healthcare Pvt Ltd ensures the highest standards in the manufacture of DAPACOOOL-GM 1 FORTE TAB, making it a trusted choice for diabetes management. For further information, contact us:

Address: 11, First Floor, Science City Road, Sola, Ahmedabad

Phone: +91 635-6666-700

Email: info@econhealthcare.net

Website: www.econhealthcare.net

Disclaimer: This product information is for general awareness only. Always take this medication under the supervision of a registered medical practitioner. Do not adjust doses or discontinue usage without consulting your doctor.

All rights reserved to Econ Healthcare