

- Sitagliptin: DPP-4 Inhibitor
- Dapagliflozin: SGLT2 Inhibitor

Indications and Uses:

- Management of type 2 diabetes to improve glycemic control.
- To be used as part of a treatment plan that includes diet and exercise.

Common Side Effects:

- Headache
- Nasal congestion
- Upper respiratory tract infection
- Back pain
- Diarrhea

Less Common Side Effects:

- Low blood sugar (hypoglycemia)
- Dehydration
- Dizziness
- Nausea
- Kidney problems

Dosage and Administration:

The recommended dose of DAPACOOOL-S TAB is one tablet once daily, taken with or without food. The dosage may vary based on the patient's condition and doctor's advice. Always follow the prescribed dose.

Warnings and Precautions:

- Monitor kidney function regularly during treatment.
- Risk of dehydration and low blood pressure, especially in elderly patients.
- May increase the risk of urinary tract infections and genital infections.
- Not recommended during pregnancy or breastfeeding.
- Take caution if you have a history of heart failure, kidney disease, or liver disorders.

Drug Interactions:

- May interact with other blood sugar-lowering medications, such as sulfonylureas or insulin, increasing the risk of hypoglycemia.
- Drugs that affect kidney function may alter the effectiveness of DAPACOOOL-S TAB.

Contraindications:

- Not for use in patients with type 1 diabetes.
- Contraindicated in patients with severe renal impairment or end-stage renal disease.
- Avoid use in patients with a history of hypersensitivity to Sitagliptin or Dapagliflozin.

Dos and Don'ts:

- Do not skip doses; take the medication as prescribed.
- Do not take more than the prescribed amount.
- Do not use this product if you are allergic to any of the ingredients.
- Do not stop taking DAPACOOOL-S TAB without consulting your doctor.
- Do not drink alcohol excessively while on this medication.

Before Taking DAPACOOOL-S TAB:

- Inform your doctor about any existing medical conditions, particularly kidney or liver issues.
- List all medications you are currently taking, including over-the-counter drugs and supplements.
- Ensure that your blood sugar levels are monitored regularly.

Disclaimer:

This information is intended for informational purposes only. The medicines should always be consumed as per the guidance and prescription of your healthcare provider. Do not alter the prescribed dosage without consulting your doctor.

All rights reserved to Econ Healthcare.