



Brand Name: ECOVOG 0.3 TAB Active Ingredient: Voglibose 0.3mg

### **General Information**:

ECOVOG 0.3 TAB contains **Voglibose**, an alpha-glucosidase inhibitor, which is used in the management of type 2 diabetes mellitus. It works by slowing down the breakdown of carbohydrates in the digestive system, thereby reducing post-meal blood sugar spikes. Voglibose is commonly used in combination with other antidiabetic medications to improve blood sugar control.

### Class:

Alpha-glucosidase Inhibitor

### Indications and Uses:

- **Type 2 Diabetes Mellitus**: ECOVOG 0.3 TAB is used to manage blood sugar levels in type 2 diabetes, especially in patients whose condition cannot be controlled by diet alone.
- **Postprandial Hyperglycemia**: It helps reduce blood sugar spikes after meals.

### Common and Less Common Side Effects:

- **Common side effects**: Gastrointestinal issues such as bloating, flatulence, diarrhea, and abdominal discomfort.
- Less common side effects: Nausea, headache, dizziness, and rash. If any of these side effects persist or worsen, consult your doctor immediately.

# Dosage and Administration:

ECOVOG 0.3 TAB is usually taken with meals, as directed by your healthcare provider. The typical dosage is **0.3mg** three times a day before meals. However, the exact dosage should be determined by your doctor based on your individual condition and response to the medication.

### Warnings and Precautions:

- **Hypoglycemia**: When taken with other antidiabetic medications, Voglibose may increase the risk of hypoglycemia (low blood sugar).
- **Gastrointestinal Issues**: Voglibose may cause bloating, flatulence, and diarrhea, especially during the initial stages of treatment.
- Liver and Kidney Impairment: Use with caution in patients with liver or kidney problems.
- **Pregnancy and Breastfeeding**: It is not recommended for use during pregnancy and breastfeeding unless advised by a doctor.

## **Drug Interactions:**

- Voglibose may interact with other antidiabetic medications, especially insulin or sulfonylureas, leading to an increased risk of hypoglycemia.
- Antacids or other gastrointestinal medications may reduce the effectiveness of Voglibose.

## Contraindications:

- Voglibose is contraindicated in patients with **hypersensitivity** to the drug or its components.
- It should not be used in patients with **severe liver or kidney impairment**, or **inflammatory bowel disease**.

## Dos and Don'ts:

- **Do** follow the prescribed dosage as directed by your healthcare provider.
- **Don't** skip meals while taking ECOVOG 0.3 TAB.
- **Do** report any unusual side effects to your doctor immediately.
- **Don't** take Voglibose with alcohol, as it may affect blood sugar levels.

### **Before Taking**:

Before starting treatment with ECOVOG 0.3 TAB, inform your doctor if you have any history of liver or kidney problems, gastrointestinal issues, or other medical conditions. Also, make sure to discuss any other medications you are currently taking to avoid interactions.

### **Quality Assurance:**

ECOVOG 0.3 TAB is a **high-quality product from Econ Healthcare Pvt Ltd**, ensuring reliable and effective management of type 2 diabetes.

For more information or inquiries, please contact us: Econ Healthcare Pvt Ltd Address: 11, First Floor, Science City Road, Sola, Ahmedabad Phone: +91 635-6666-700 Email: info@econhealthcare.net Website: <u>www.econhealthcare.net</u>

**Disclaimer**: This information is provided for general knowledge and is not intended to replace the guidance of a healthcare provider. Always follow your doctor's advice regarding dosage and use.

# All rights reserved to Econ Healthcare.