



<p>Rx Febuxostat Tablets Febstar[®]-40 Tablets फेबस्टार-४०</p> <p>10 x 10 Tablets</p>	
---	--

Composition :
Each film coated tablet contains :
Febuxostat 40 mg
Excipients Q.S.
Color : Titanium Dioxide IP
Dose: As directed by the Physician
Storage :
Store protected from light & moisture
at a temperature not exceeding 30° C.

Febstar[®] - 40
फेबस्टार-४० Tablets
Keep medicine out of reach of children

CAUTION : Not to be sold by retail
without the prescription of a
Registered Medical Practitioner.

Mfg. by :Shantam Pharmaceuticals Pvt. Ltd.
546/2, At. Rakanpur, Ta. Kalol,Dist.Gandhinagar - 382 721
www.shantampharmaceuticals.com

Brand Name: FEBSTAR-40 TAB

Composition: Febuxostat 40mg

General Information

FEBSTAR-40 TAB contains Febuxostat, a xanthine oxidase inhibitor used to manage hyperuricemia in adults with gout. By decreasing uric acid production in the body, it helps prevent gout attacks and complications caused by high uric acid levels.

Class:

Xanthine Oxidase Inhibitor

Indications and Uses:

- Treatment of chronic hyperuricemia in patients with gout.
- Management of uric acid levels in conditions causing overproduction of uric acid.

Common and Less Common Side Effects:

Common Side Effects:

- Nausea
- Joint pain
- Rash
- Liver enzyme abnormalities

Less Common Side Effects:

- Dizziness
- Fatigue
- Diarrhea
- Cardiovascular symptoms like chest pain (rare)

Doses and Administration:

The usual recommended dose is one tablet (40mg) once daily, with or without food. Dosage may vary based on the patient's condition and doctor's advice.

Warnings and Precautions:

- Regular monitoring of uric acid levels and liver function tests is advised.
- Patients with cardiovascular disease should use with caution.
- Avoid using during pregnancy or breastfeeding unless advised by a doctor.

Drug Interactions:

FEBSTAR-40 may interact with:

- Azathioprine
 - Mercaptopurine
 - Theophylline
- Inform your doctor about all medications you are currently taking to avoid potential interactions.

Contraindications:

- Hypersensitivity to Febuxostat or any component of the medication.
- Severe liver or kidney impairment.

Dos and Don'ts:**Dos:**

- Take medication as prescribed by your doctor.
- Maintain a low-purine diet to support treatment.
- Stay hydrated to help prevent kidney stones.

Don'ts:

- Do not stop taking the medication without consulting your doctor.
- Avoid alcohol as it can increase uric acid levels.
- Do not take with other medications without prior medical consultation.

Before Taking FEBSTAR-40 TAB:

- Inform your doctor if you have a history of liver or kidney disease.

- Discuss any history of cardiovascular problems or allergies.

A Quality Product from Econ Healthcare

Address: 11, First Floor, Science City Road, Sola, Ahmedabad

Phone: +91 635-6666-700

Email: info@econhealthcare.net

Website: www.econhealthcare.net

Disclaimer: This information is for general awareness and is not a substitute for professional medical advice. Always follow your doctor's instructions regarding medication use. Do not modify or discontinue doses prescribed by your doctor.

All rights reserved to Econ Healthcare.