



Rx
Meropenem Injection IP

Merocon
For IV USE ONLY
Single Dose Vial

(Combipack with Sterile Water for Injections IP)

1 g

Each combipack contains:
(a) 1 Vial of Meropenem Injection IP
Each vial contains :
Meropenem IP (Sterile) eq. to
Anhydrous Meropenem 1000 mg
Sodium
(as Sodium Carbonate) 90.2 mg
(b) 1 Ampoule of Sterile Water for Injections IP
Each ampoule contains:
Sterile Water for Injections IP 20 ml
Dosage: As directed by the Physician.
Store below 30°C. Protect from moisture & light. Do not freeze.
Direction for use:
Dissolve the contents of vial in 20 ml of Sterile Water for Injections IP for IV use. Use freshly prepared solution, effervescence occurs on addition of Sterile Water for Injections IP.
The constituted solution should be used immediately after preparation.
Caution: If any foreign matter is visible in the vial after dissolving the contents, please do not use the solution.
Keep medicine out of reach of children.

SCHEDULE H1 PRESCRIPTION DRUG-CAUTION
It is dangerous to take this preparation except in accordance with the medical advice.
Not to be sold by retail without the prescription of a Registered Medical Practitioner.

Manufactured by:
Akums Drugs & Pharmaceuticals Ltd.
2,3,4 & 5, Sector-6B, I.I.E., SIDCUL,
Ranipur, Haridwar-249 403, INDIA.

MEROCON-1GM INJ

Composition: Each vial contains **Meropenem Injection 1gm IP**

General Information

MEROCON-1GM INJ is a broad-spectrum antibiotic belonging to the **carbapenem class** of drugs. Meropenem is used to treat severe bacterial infections caused by susceptible organisms. It works by interfering with the bacterial cell wall synthesis, leading to the destruction of the bacteria.

Indications and Uses

- Severe infections of the lungs, urinary tract, skin, and soft tissues.
 - Intra-abdominal infections.
 - Bacterial meningitis.
 - Septicemia.
-

Common and Less Common Side Effects

Common side effects:

- Nausea and vomiting.
- Diarrhea.
- Injection site reactions such as redness or swelling.

Less common side effects:

- Headache.
 - Dizziness.
 - Rash or allergic reactions.
-

Doses and Administration

- Administered as an intravenous (IV) injection or infusion.
 - The dosage and duration depend on the severity of the infection and the patient's condition.
 - Should only be used under the guidance of a healthcare professional.
-

Warnings and Precautions

- Use with caution in patients with kidney impairment.
 - Inform the doctor of any history of allergic reactions to antibiotics.
 - Do not self-administer this injection.
-

Drug Interactions

- May interact with probenecid, valproic acid, or other antibiotics.
 - Inform the doctor about all ongoing medications.
-

Contraindications

- Not recommended for patients with a known allergy to meropenem or other beta-lactam antibiotics.
-

Dos and Don'ts

Dos:

- Follow the doctor's prescription carefully.
- Monitor for any allergic reactions post-administration.

Don'ts:

- Do not administer without medical supervision.
 - Avoid using in patients with severe allergic reactions to penicillins or cephalosporins.
-

Before Taking MEROCON-1GM INJ

- Inform your doctor about any pre-existing liver or kidney conditions.
 - Share your complete medical history, including ongoing medications.
 - Avoid use during pregnancy or breastfeeding unless advised by your doctor.
-

A Quality Product from Econ Healthcare

Econ Healthcare Pvt Ltd is committed to delivering high-quality healthcare products.

Address: **11, First Floor, Science City Road, Sola, Ahmedabad**

Phone: **+91 635-6666-700**

Email: **info@econhealthcare.net**

Website: www.econhealthcare.net

Disclaimer

This is general information provided for better understanding of the product. Always follow your doctor's guidance regarding the use of this medication. Do not change or stop the prescribed dose without consulting your doctor.

All rights reserved to Econ Healthcare Pvt Ltd.